

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

site, but this point will be investigated further before any definite recommendation is made as to the location of the substations. It is purposed to continue the work at the main station throughout at least one sun-spot period, and for two or three summers at the two substations.

With the foresight and thoroughness with which the committee has made its plans and preliminary investigations as an earnest of the wise future administration of its trust, with the powerful equipment for which these recommendations call, and with plans founded on the notable advances of the past twenty years, astrophysicists may well hope for a decade of unparalleled progress in the field of solar physics.

## PLANETARY PHENOMENA FOR JULY AND AUGUST, 1904.

## By Malcolm McNeill.

## PHASES OF THE MOON, PACIFIC TIME.

```
Last Quarter, July 5, 8h 54m P.M. Last Quarter, Aug. 4, 6h 3m A.M. New Moon, '' 12, 9 27 P.M. New Moon, '' 11, 4 58 A.M. First Quarter, '' 19, 12 49 P.M. First Quarter, '' 17, 8 27 P.M. Full Moon, '' 25, 5 2 P.M.
```

The Earth is in aphelion July 4th, 5 P.M., Pacific time.

Mercury is a morning star on July 1st, too near the Sun to be seen, rising about three quarters of an hour before sunrise. It rapidly approaches the Sun, passes superior conjunction, and becomes an evening star on July 9th. By August 1st it has receded far enough from the Sun to set rather more than an hour after sunset, and this interval does not change very much until after the middle of the month. It reaches greatest east elongation on the evening of August 19th. Its apparent distance from the Sun will then be 27° 24'. This elongation is much greater than the average greatest elongation, since the planet passed its aphelion on August 17th, only two days before. After passing greatest elongation the planet approaches

the Sun quite rapidly, and at the end of the month it sets about half an hour after sunset. It is in close conjunction with *Mars* on the morning of July 2d, but both bodies are almost too near the Sun for observation at that time. Also, it is in conjunction with *Venus* only ten hours after passing conjunction with the Sun.

Venus is a morning star too near the Sun to be easily seen on July 1st. It comes to superior conjunction July 7th, 11 p.m., Pacific time. It then becomes an evening star, and will remain so until the following spring. Its apparent distance from the Sun increases rather slowly, and its path among the stars tends toward the south; so the interval between the setting of the Sun and of the planet does not grow very rapidly. On August 1st it is only thirteen minutes, and on August 31st about forty minutes. Notwithstanding its brightness, now at its minimum, it will not be easy to see it with the naked eye until nearly the close of August. It is in perihelion on July 23d, but its orbit is so nearly circular that the difference between greatest and least distance from the Sun is less than half a million miles.

Mars is now getting far enough away from the Sun to be seen as a morning star. On July 1st it rises about forty minutes before sunrise, on August 1st an hour and a half before, and on August 31st well over two hours before. During the two months it moves over 40° eastward and 6° southward in Gemini. During the middle and latter part of August it is a few degrees south of Castor and Pollux, a and B Geminorum.

Jupiter rises half an hour after midnight on July 1st, and by the end of August at a little before 9 P.M. It moves eastward in the eastern part of *Pisces* until August 20th, about 3°, and then begins to move slowly westward.

Saturn rises before 10 P.M. on July 1st, and at about 5:40 on August 30th. So it is in fair position for early-evening observation in August. It comes to opposition with the Sun on August 10th, and is then above the horizon during the entire night. During the two months it moves about  $4^{\circ}$  westward in the constellation Capricorn, somewhat east of  $\alpha$  and  $\beta$ , the principal stars of the constellation. The rings apparently widen out a trifle from their minimum in June, on account of the Earth's motion in reference to their plane; but aside

from this temporary increase in apparent breadth, there will be a general progressive diminution until the motion of the planet brings the plane of the rings into coincidence with the Earth.

Uranus crosses the meridian shortly after II P.M., on July Ist, and at 7 P.M. on September Ist. So it is in fair position for evening observation. It moves a little less than 2° westward in the western part of Sagittarius, becoming stationary early in September. No bright star is near enough to make the planet's identification easy.

Neptune is a morning object in Gemini. By the end of August it rises not long after midnight.

It will be worth while to look for the annual *Perseid* meteors during the second week in August. The meteors are so scattered that we get some every year, while there is no great shower occurring at intervals of several years,—or failing to come to time in anything like its hoped for brilliancy, as happened with the *Leonids* a few years ago.